Available online at www.sciencedirect.com
ScienceDirect
Nuclear Physics B Proceedings Supplement 00 (2014) 1–3
Nuclear Physics B Proceedings Supplement

Hadronic total cross sections, Wilson loop correlators and the QCD spectrum

Matteo Giordano[a,1] and Enrico Meggiolaro[b]

[a]Institute for Nuclear Research of the Hungarian Academy of Sciences (ATOMKI), Bem tér 18/c, H–4026 Debrecen, Hungary
[b]Dipartimento di Fisica, Università di Pisa, and INFN, Sezione di Pisa, Largo Pontecorvo 3, I–56127 Pisa, Italy

arXiv:1409.7376v1 [hep-ph] 25 Sep 2014

Abstract

We show how to obtain rising hadronic total cross sections in QCD, in the framework of the nonperturbative approach to soft high-energy scattering based on Wilson-loop correlators. Total cross sections turn out to be of "Froissart"-type, i.e., the leading energy dependence is of the form $\sigma_{\rm tot} \sim B \log^2 s$, in agreement with experiments. The observed universality of the prefactor B is obtained rather naturally in this framework. In this case, B is entirely determined by the stable spectrum of QCD, and predicted to be $B_{th} = 0.22\,\text{mb}$, in fair agreement with experiments.

Keywords: total cross sections, QCD, nonperturbative approach

1. Introduction

Explaining the behaviour of hadronic total cross sections at high energy is a very old problem, which is very rarely attacked within the framework of QCD. Most of the approaches to this problem are based on phenomenological models, which are sometimes QCD-inspired, but a full derivation from first principles of QCD is still lacking.

The observed rise of hadronic total cross sections at high energy is well described by a "Froissart-like" behaviour $\sigma_{tot} \sim B \log^2 s$ with a universal prefactor $B \simeq 0.27 \div 0.28\,\text{mb}$ [1]. This behaviour respects unitarity, as encoded in the Froissart-Łukaszuk-Martin bound [2, 3, 4], $\sigma_{tot} \leq B_{\rm FLM} \log^2 \frac{s}{s_0}$, since $B \ll B_{\rm FLM} = \pi/m_\pi^2 \simeq 65\,\text{mb}$.

Besides deriving σ_{tot} from the first principles of QCD, one should be able to explain the observed universality of B, and also the two orders of magnitude separating B and $B_{\rm FLM}$. A step in this direction has been made in Ref. [5], where we have derived the asymptotic behaviour of hadronic total cross sections in the framework of the nonperturbative approach to soft high-energy scattering [6, 7, 8], finding indeed a "Froissart-like" behaviour, and a theoretical prediction for B in fair agreement with experiments.

2. Nonperturbative approach to soft high-energy scattering

Understanding the rise of σ_{tot} is part of the problem of soft high-energy scattering, characterised by $|t| \leq 1\,\text{GeV}^2 \ll s$. In this regime perturbation theory is not fully reliable, and a nonperturbative approach is needed. In a nutshell, the nonperturbative approach to *elastic meson-meson scattering* is as follows [6, 7, 8]:

1. mesons are described as wave packets of transverse colourless dipoles;
2. in the soft high-energy regime, the dipoles travel essentially undisturbed on their classical, almost lightlike trajectories;
3. mesonic amplitudes are obtained from the dipole-dipole (dd) amplitudes after folding with the appropriate squared wave functions, $A(s,b) = \langle\!\langle A^{(dd)}(s,b;\nu)\rangle\!\rangle$.

Email addresses: giordano@atomki.mta.hu (Matteo Giordano), enrico.meggiolaro@df.unipi.it (Enrico Meggiolaro)

[1]Supported by the Hungarian Academy of Sciences under "Lendület" grant No. LP2011-011.



Figure 1: The relevant Euclidean Wilson loops.

Here b is the impact parameter[2], ν denotes collectively the dipole variables (longitudinal momentum fraction, transverse size and orientation), and $\langle\!\langle \ldots \rangle\!\rangle$ stands for integration over the dipole variables with the mesonic wave functions. This approach extends also to processes involving baryons, if one adopts for them a quark-diquark picture [9].

At high energy the dd amplitude is given by the normalised connected correlator C_M of the Wilson loops (WL) running along the classical trajectories of the two dipoles, $A^{(dd)}(s,b;\nu) = -C_M(\chi,b;\nu)$, with $\chi \simeq \log\frac{s}{m^2}$ the hyperbolic angle between the trajectories, and m the mass of the mesons (taken to be equal for simplicity).

The correlator C_M is obtained from the correlator C_E of two Euclidean WL at angle θ (see Fig. 1),

$$C_E(\theta,b;\nu) \equiv \lim_{T\to\infty} \frac{\langle \mathcal{W}_{C_1}\mathcal{W}_{C_2}\rangle}{\langle \mathcal{W}_{C_1}\rangle\langle \mathcal{W}_{C_2}\rangle} - 1\,, \tag{1}$$

through the analytic continuation (AC) [10, 11]

$$C_M(\chi,b;\nu) = C_E(\theta\to -i\chi,b;\nu)\,. \tag{2}$$

A more detailed discussion of the approach can be found in Ref. [5] and references therein. The Euclidean formulation has allowed the study of the relevant WL correlator by means of nonperturbative techniques, which include instantons [12, 13], the model of the stochastic vacuum [14], holography [15, 16, 17, 18, 19], and the lattice [13, 20, 21].

[2]We are interested in unpolarised scattering, so only the impact parameter modulus matters; for definiteness, we keep it oriented along direction 2 in the transverse plane.

3. Large-s behaviour of $\sigma_{\rm tot}$

The energy dependence of $\sigma_{\rm tot}$ is determined by the large-s, large-b behaviour of the amplitude through the "effective radius" of interaction $b_c = b_c(s)$, beyond which the amplitude is negligible, as $\sigma_{\rm tot} \propto b_c^2$.

To determine b_c, in Ref. [5] we employed the following strategy. On the Euclidean side, we obtain information on the b- and θ-dependencies of C_E by inserting between the WL operators (for large enough b) a complete set of asymptotic states, characterised by their particle content and by the momenta and spin of each particle. After AC this gives us information on the s- and b-dependencies. This requires two crucial analyticity assumptions:

1. AC can be performed separately for each term in the sum;
2. WL matrix elements are analytic in θ.

A few reasonable finiteness assumptions on the WL matrix elements are also made.

Under these assumptions, at large χ, b, the relevant Minkowskian correlator reads [5]

$$C_M(\chi,b;\nu) \simeq \sum_{\alpha\neq 0} f_\alpha(\nu) \prod_a [w_a(\chi,b)]^{n_a(\alpha)}\,, \tag{3}$$

where the sum is over (non-vacuum) states α, and $n_a(\alpha)$ is the number of particles of type a in state α. Here

$$w_a(\chi,b) = \frac{e^{[r^{(a)}\chi - b]m^{(a)}}}{\sqrt{2\pi b m^{(a)}}}\,, \qquad r^{(a)} \equiv \frac{s^{(a)}-1}{m^{(a)}}\,, \tag{4}$$

with $(s^{(a)}, m^{(a)})$ spin and mass of particles of type a, and f_α are functions of the dipole variables only. Particles of type a contribute only for $b \lesssim r^{(a)}\chi$, and so the effective radius of interaction is given by

$$b_c(s) = \left(\max_a r^{(a)}\right)\log\frac{s}{m^2} \equiv \frac{1}{\mu}\log\frac{s}{m^2}\,. \tag{5}$$

Here we assume the maximum to exist and to be positive. If it were zero or negative, $\sigma_{\rm tot}$ would be constant or vanishing at high energy. The spectrum is supposed to be free of massless states: in case they were present and with spin at most 1, the maximisation should be performed on the massive spectrum only [5].

Using Eq. 3 we find for $\sigma_{\rm tot}$

$$\sigma_{\rm tot} \underset{s\to\infty}{\simeq} 2\pi(1-\kappa)[b_c(s)]^2 \simeq \frac{2\pi}{\mu^2}(1-\kappa)\log^2\frac{s}{m^2}\,, \tag{6}$$

with $|\kappa| \leq 1$ due to unitarity [5]. In general κ depends on the colliding hadrons. Analyticity and crossing symmetry [22, 23] requirements show that universality is most naturally achieved if $\kappa = 0$, corresponding to a vanishing or oscillating correlator as $\chi \to \infty$ at fixed b.



Figure 2: Plot of $2\pi\left(\frac{s-1}{m}\right)^2$ for QCD-stable states. The 2^{++} glueball state [24, 25] and the experimental value for B [1] are also shown. Nuclear data are taken from Ref. [26].

4. Total cross sections from the hadronic spectrum

The total cross section satisfies the bound [5]

$$\sigma_{\text{tot}} \underset{s\to\infty}{\lesssim} \frac{4\pi}{\mu^2}\log^2\frac{s}{m^2} = 2B_{\text{th}}\log^2\frac{s}{m^2}\,, \tag{7}$$

with $\mu^{-1} = \max_a r^{(a)}$ determined from the hadronic spectrum by maximising $r^{(a)}$ over the stable states of QCD *in isolation*, as electroweak effects have been neglected from the onset. Only states with $s^{(a)} \geq 1$ are considered, including nuclei (see Fig. 2).

Quite surprisingly, this singles out the $\Omega^{\pm}$ baryon, which yields $B_{\text{th}} \simeq 0.56$ GeV^{-2}. The resulting bound on σ_{tot} is much more stringent than the original bound, and the prefactor is of the same order of magnitude of the experimental value $B_{\text{exp}} \simeq 0.69 \div 0.73$ GeV^{-2} [1]. Furthermore, our bound is not singular in the chiral limit [27].

If universality is achieved as discussed above in Section 3, then σ_{tot} is entirely determined by the hadronic spectrum, and reads [5]

$$\sigma_{\text{tot}} \underset{s\to\infty}{\simeq} \frac{2\pi}{\mu^2}\log^2\frac{s}{m^2} = B_{\text{th}}\log^2\frac{s}{m^2}\,. \tag{8}$$

This *prediction* for the prefactor is in fair agreement with B_{exp}, taking into account a systematic error of order 10% on B_{exp}, estimated by comparing the results of different fitting procedures [1, 28, 29, 30]. The same conditions leading to universal total cross sections also give universal, black-disk-like elastic scattering amplitudes [5].

Total cross sections are usually believed to be governed by the gluonic sector of QCD. However, in the *quenched* theory one finds from the glueball spectrum $B_Q \gtrsim 1.6 B_{\text{exp}}$, suggesting large unquenching effects [5].

5. Conclusions

In Ref. [5] we have derived the asymptotic, high-energy behaviour of hadronic total cross sections in the framework of the nonperturbative approach to soft high-energy scattering [6, 7, 8]. We find a "Froissart-like" behaviour $\sigma_{tot} \sim B\log^2 s$, with B (mainly) determined by the hadronic spectrum (see Eqs. 7 and 8), and in fair agreement with experiments.

Our main results do not depend on the detailed description of the hadrons in terms of partons: adding gluons and sea quarks to the wave functions would lead to more complicated WLs, but since our argument is independent of their detailed form, our conclusions remain unchanged.

References

[1] K. A. Olive *et al.* [Particle Data Group Collaboration], Chin. Phys. C **38** (2014) 090001.
[2] M. Froissart, Phys. Rev. **123** (1961) 1053.
[3] A. Martin, Il Nuovo Cimento **42A** (1966) 930.
[4] L. Łukaszuk and A. Martin, Il Nuovo Cimento **52A** (1967) 122.
[5] M. Giordano and E. Meggiolaro, JHEP **1403** (2014) 002.
[6] O. Nachtmann, Ann. Phys. **209** (1991) 436.
[7] H. G. Dosch, E. Ferreira and A. Krämer, Phys. Rev. D **50** (1994) 1992.
[8] S. Donnachie, G. Dosch, P. Landshoff and O. Nachtmann, *Pomeron Physics and QCD* (Cambridge University Press, Cambridge, 2002).
[9] M. Rueter and H. G. Dosch, Phys. Lett. B **380** (1996) 177.
[10] E. Meggiolaro, Nucl. Phys. B **707** (2005) 199.
[11] M. Giordano and E. Meggiolaro, Phys. Lett. B **675** (2009) 123.
[12] E. Shuryak and I. Zahed, Phys. Rev. D **62** (2000) 085014.
[13] M. Giordano and E. Meggiolaro, Phys. Rev. D **81** (2010) 074022.
[14] A. I. Shoshi, F. D. Steffen, H. G. Dosch and H. J. Pirner, Phys. Rev. D **68** (2003) 074004.
[15] R. A. Janik and R. Peschanski, Nucl. Phys. B **565** (2000) 193.
[16] R. A. Janik and R. Peschanski, Nucl. Phys. B **586** (2000) 163.
[17] R. A. Janik, Phys. Lett. B **500** (2001) 118.
[18] M. Giordano and R. Peschanski, JHEP **1005** (2010) 037.
[19] G. Başar, D. E. Kharzeev, H.-U. Yee, and I. Zahed, Phys. Rev. D **85** (2012) 105005.
[20] M. Giordano and E. Meggiolaro, Phys. Rev. D **78** (2008) 074510.
[21] M. Giordano, E. Meggiolaro and N. Moretti, JHEP **1209** (2012) 031.
[22] M. Giordano and E. Meggiolaro, Phys. Rev. D **74** (2006) 016003.
[23] E. Meggiolaro, Phys. Lett. B **651** (2007) 177.
[24] C. J. Morningstar and M. J. Peardon, Phys. Rev. D **60** (1999) 034509.
[25] E. Gregory *et al.*, JHEP **1210** (2012) 170.
[26] G. Audi, O. Bersillon, J. Blachot and A. H. Wapstra, Nucl. Phys. A **729** (2003) 3.
[27] M. Giordano and E. Meggiolaro, in preparation.
[28] J. R. Cudell *et al.* [COMPETE collaboration], Phys. Rev. D **65** (2002) 074024.
[29] M. Ishida and K. Igi, Phys. Lett. B **670** (2009) 395.
[30] M. M. Block and F. Halzen, Phys. Rev. Lett. **107** (2011) 212002.